



**MEDICATION DEFERRAL LIST**

**DO NOT STOP taking medications prescribed by your doctor in order to donate blood.**

Donating while taking these drugs could have a negative effect on your health or on the health of the recipient of your blood.

**PLEASE TELL US IF YOU:**

ARE BEING TREATED WITH ANY OF THE FOLLOWING TYPES OF MEDICATIONS:	OR HAVE TAKEN:	WHICH IS ALSO CALLED:		ANYTIME IN THE LAST:
Antiplatelet agents (usually taken to prevent stroke or heart attack)	Feldene	piroxicam		2 days
	Effient	prasugrel		3 days
	Brilinta	ticagrelor		7 days
	Plavix	clopidogrel		14 days
	Ticlid	ticlopidine		
	Zontivity	vorapaxar		1 month
Anticoagulants or "blood thinners" (usually taken to prevent blood clots in the legs and lungs and to prevent strokes)	Arixtra	fondaparinux		2 days
	Eliquis	apixaban		
	Fragmin	dalteparin		
	Lovenox	enoxaparin		
	Pradaxa	dabigatran		
	Savaysa	edoxaban		
	Xarelto	rivaroxaban		
	Coumadin, Warfilone, Jantoven	warfarin		7 days
Acne treatment	Heparin, low molecular weight heparin			1 Month
	Accutane	isotretinoin		
	Amnesteem			
	Absorica			
	Claravis			
	Myorisan			
	Sotret			
Zenatane				
Multiple myeloma	Thalomid	Thalidomide		6 Months
Rheumatoid arthritis	<b>Revlimid</b>	<b>lenalidomide</b>		
Hair loss remedy	Rinvoq	upadacitinib		
Prostate symptoms	Propecia	finasteride		
	Proscar	finasteride		
	Avodart	dutasteride		
	Jalyn			
Immunosuppressant	Cellcept	mycophenolate mofetil		6 weeks
Hepatitis exposure	Hepatitis B Immune Globulin	HBIG		3 months
HIV Prevention (also known as PrEP or PEP)	Any medication taken by mouth (oral) to prevent HIV	Truvada	emtricitabine and tenofovir disoproxil fumarate	
		Descovy	emtricitabine and tenofovir alafenamide	
	<b>Injectable HIV prevention</b>	<b>Apretude</b>	<b>cabotegravir</b>	2 Years
Basal cell skin cancer	Erivedge	vismodegib sonidegib		2 years
	Odomzo			
Relapsing multiple sclerosis	Aubagio	teriflunomide		
Rheumatoid arthritis	Arava	leflunomide		3 Years
Psoriasis	Soriatane	acitretin		
	Tegison	etretinate		
HIV treatment	<b>Any medication to treat HIV. May also be called antiretroviral therapy (ART)</b>			Ever
Experimental Medication	<b>Any experimental medications used for any condition</b>			12 Months

For additional information, please see the back page.

Facility Name: Sylvan N. Goldman Oklahoma Blood Institute 1001 N. Lincoln Blvd., Oklahoma City, Oklahoma 73104. The official copy of blood bank documentation is the electronic copy on file with the local area network. The official copy of records created from forms is paper unless designated otherwise.

Apply Location sticker here OR Insert Location code _____
---



**MEDICATION DEFERRAL LIST**

**DO NOT STOP taking medications prescribed by your doctor in order to donate blood.**

*Some medications affect your eligibility as a blood donor for the following reasons:*

**Antiplatelet agents affect platelet function**, so people taking these drugs should not donate platelets for the indicated time. You may still be able to donate whole blood or red blood cells by apheresis.

**Anticoagulants or "blood thinners"** are used to treat or prevent blood clots in the legs, lungs, or other parts of the body, and to prevent strokes. These medications affect the blood's ability to clot, which might cause excessive bruising or bleeding when you donate. You may still be able to donate whole blood or red blood cells by apheresis.

**Isotretinoin, finasteride, dutasteride, acitretin, and etretinate** can cause birth defects. Your donated blood could contain high enough levels to damage the unborn baby if transfused to a pregnant woman.

**Thalomid (thalidomide), Revlimid (lenalidomide), Erivedge (vismodegib), Odomxo (sonidegib), Aubagio (teriflunomide), and Rinvoq (upadacitinib)** may cause birth defects or the death of an unborn baby if transfused to a pregnant woman.

**Cellcept (mycophenolate mofetil) and Arava (leflunomide)** are immunosuppressants which may cause birth defects or the death of an unborn baby if transfused to a pregnant woman.

**PrEP or pre-exposure prophylaxis** involves taking a specific combination of oral medicines (i.e., short-acting antiviral PrEP) or injections (i.e., long-acting antiviral PrEP) as a prevention method for people who are HIV negative and at high risk of HIV infection. FDA has determined that the available data demonstrate that the use of PrEP or PEP may delay the detection of HIV by currently licensed screening tests for blood donations, potentially resulting in false negative results in infected individuals. Although "Undetectable = Untransmittable" for sexual transmission, this **does not apply to transfusion transmission**.

**PEP or post-exposure prophylaxis** is a short-acting treatment started as soon as possible after a high-risk exposure to HIV to reduce the risk of infection. FDA has determined that the available data demonstrate that the use of PrEP or PEP may delay the detection of HIV by currently licensed screening tests for blood donations, potentially resulting in false negative results in infected individuals. Although "Undetectable = Untransmittable" for sexual transmission, this **does not apply to transfusion transmission**.

**ART or antiretroviral therapy** is the use of a combination of HIV medicines (called an HIV regimen) to treat HIV infection. HIV infection requires a permanent deferral despite treatment with ART. Antiretroviral drugs do not fully eliminate the virus from the body, and donated blood from individuals infected with HIV taking ART can potentially still transmit HIV to a transfusion recipient. Although "Undetectable = Untransmittable" for sexual transmission, this **does not apply to transfusion transmission**.

**Hepatitis B Immune Globulin (HBIG)** is an injected material used to prevent hepatitis B infection following a possible or known exposure to hepatitis B. HBIG does not prevent hepatitis B infection in every case, therefore, persons who have received HBIG must wait to donate blood.

**Experimental Medication or Unlicensed (Experimental) Vaccine** is usually associated with a research study, and the effect on the safety of transfused blood is unknown.

ADDITIONAL MEDICATIONS PLEASE TELL US IF YOU:			
Are being treated with the following types of medications:	or have taken:	which is also called:	anytime in the last:
<b>Antiplatelet agents</b>	Persantine	dipyridamole	2 days
	Aggrenox	aspirin/extended-release dipyridamole	
	Fiorinal Fiormor Fiortal Fortabs Lanroif	butalbital/aspirin/caffeine	
	Soma Compound	aspirin and carisoprodol	
<b>Prostate Cancer</b>	Lupron Lupron Depot Eligrard Viadur	leuprolide acetate	6 months
<b>Menstrual disorder treatment Or Endometriosis</b>	Aygestin	norethindrone acetate  (If you are taking norethindrone acetate for birth control ONLY, you are eligible to donate today.)	
<b>Atrial fibrillation treatment</b>	Multaq	dronedarone	
<b>Direct thrombin inhibitors</b>	Acova	argatroban	2 days
<b>Direct Xa inhibitors</b>	Betrixaban Darexaban Otamixaban	N/A	
		Lixiana	edoxaban
<b>Acne treatment</b>	Tazorac cream	tazarotene	1 month
<b>Autoimmune treatment</b>	IVIG	intravenous immunoglobulin	12 months

Facility Name: Sylvan N. Goldman Oklahoma Blood Institute 1001 N. Lincoln Blvd., Oklahoma City, Oklahoma 73104. The official copy of blood bank documentation is the electronic copy on file with the local area network. The official copy of records created from forms is paper unless designated otherwise.

Apply Location sticker here  
OR  
Insert Location code \_\_\_\_\_